

Table 1 DISCRETE DISTRIBUTIONS

Name of parametric family of distributions	Discrete density functions $f(x)$	Parameter space	Mean $\mu = E[X]$
Discrete uniform	$f(x) = \frac{1}{N} I_{(1, \dots, N)}(x)$	$N = 1, 2, \dots$	$\frac{N+1}{2}$
Bernoulli	$f(x) = p^x q^{1-x} I_{(0, 1)}(x)$	$0 \leq p \leq 1$ ( $q = 1 - p$ )	$p$
Binomial	$f(x) = \binom{n}{x} p^x q^{n-x} I_{(0, 1, \dots, n)}(x)$	$0 \leq p \leq 1$ $n = 1, 2, 3, \dots$ ( $q = 1 - p$ )	$np$
Hypergeometric	$f(x) = \frac{\binom{K}{x} \binom{M-K}{n-x}}{\binom{M}{n}} I_{(0, 1, \dots, n)}(x)$	$M = 1, 2, \dots$ $K = 0, 1, \dots, M$ $n = 1, 2, \dots, M$	$\frac{K}{M}$
Poisson	$f(x) = \frac{e^{-\lambda} \lambda^x}{x!} I_{(0, 1, \dots)}(x)$	$\lambda > 0$	$\lambda$
Geometric	$f(x) = pq^x I_{(0, 1, \dots)}(x)$	$0 < p \leq 1$ ( $q = 1 - p$ )	$\frac{q}{p}$
Negative binomial	$f(x) = \binom{r+x-1}{x} p^x q^{r-x} I_{(0, 1, \dots)}(x)$	$0 < p \leq 1$ $r > 0$ ( $q = 1 - p$ )	$\frac{rq}{p}$

Variance $\sigma^2 = E[(X - \mu)^2]$	Moments $\mu_r^* = E[X^r]$ or $\mu_r = E[(X - \mu)^r]$ and/or cumulants $\kappa_r$	Moment generating function $E[e^{tX}]$
$\frac{N^2 - 1}{12}$	$\mu_3^* = \frac{N(N+1)^2}{4}$ $\mu_4^* = \frac{(N+1)(2N+1)(3N^2+3N-1)}{30}$	$\sum_{j=1}^N \frac{1}{N} e^{jt}$
$pq$	$\mu_r^* = p$ for all $r$	$q + pe^t$
$npq$	$\mu_3 = npq(q-p)$ $\mu_4 = 3n^2 p^2 q^2 + npq(1-6pq)$	$(q + pe^t)^n$
$\frac{K}{n} \frac{M-K}{M} \frac{M-n}{M-1}$	$E[X(X-1) \cdots (X-r+1)] = r! \frac{\binom{K}{r} \binom{M-n}{r}}{\binom{M}{r}}$	not useful
$\lambda$	$\kappa_r = \lambda$ for $r = 1, 2, \dots$ $\mu_3 = \lambda$ $\mu_4 = \lambda + 3\lambda^2$	$\exp[\lambda(e^t - 1)]$
$\frac{q}{p^2}$	$\mu_3 = \frac{q+q^2}{p^2}$ $\mu_4 = \frac{q+7q^2+q^3}{p^3}$	$\frac{p}{1-qe^t}$
$\frac{rq}{p^2}$	$\mu_3 = \frac{r(q+q^2)}{p^2}$ $\mu_4 = \frac{r[q+(3r+4)q^2+q^3]}{p^3}$	$\left(\frac{p}{1-qe^t}\right)^r$

Table 2. CONTINUOUS DISTRIBUTIONS

Name of parametric family of distributions	Cumulative distribution function $F(\cdot)$ or probability density function $f(\cdot)$	Parameter space	Mean $\mu = \mathcal{E}[X]$
Uniform or rectangular	$f(x) = \frac{1}{b-a} I_{(a,b)}(x)$	$-\infty < a < b < \infty$	$\frac{a+b}{2}$
Normal	$f(x) = \frac{1}{\sqrt{2\pi\sigma}} \exp[-(x-\mu)^2/2\sigma^2]$	$-\infty < \mu < \infty$ $\sigma > 0$	$\mu$
Exponential	$f(x) = \lambda e^{-\lambda x} I_{(0,\infty)}(x)$	$\lambda > 0$	$\frac{1}{\lambda}$
Gamma	$f(x) = \frac{\lambda^r}{\Gamma(r)} x^{r-1} e^{-\lambda x} I_{(0,\infty)}(x)$	$\lambda > 0$ $r > 0$	$\frac{r}{\lambda}$
Beta	$f(x) = \frac{1}{B(a,b)} x^{a-1} (1-x)^{b-1} I_{(0,1)}(x)$	$a > 0$ $b > 0$	$\frac{a}{a+b}$
Cauchy	$f(x) = \frac{1}{\pi\beta} \frac{1}{(1 + ((x-\alpha)/\beta)^2)}$	$-\infty < \alpha < \infty$ $\beta > 0$	Does not exist
Lognormal	$f(x) = \frac{1}{x\sqrt{2\pi\sigma}} \exp[-(\log_e x - \mu)^2/2\sigma^2] I_{(0,\infty)}(x)$	$-\infty < \mu < \infty$ $\sigma > 0$	$\exp[\mu + \frac{1}{2}\sigma^2]$
Double exponential	$f(x) = \frac{1}{2\beta} \exp\left(-\frac{ x-\alpha }{\beta}\right)$	$-\infty < \alpha < \infty$ $\beta > 0$	$\alpha$

Variance $\sigma^2 = \mathcal{E}[(X-\mu)^2]$	Moments $\mu_r' = \mathcal{E}[X^r]$ or $\mu_r = \mathcal{E}[(X-\mu)^r]$ and/or cumulants $\kappa_r$	Moment generating function $\mathcal{E}[e^{tX}]$
$\frac{(b-a)^2}{12}$	$\mu_r = 0$ for $r$ odd $\mu_r = \frac{(b-a)^r}{2^r(r+1)}$ for $r$ even	$\frac{e^{bt} - e^{at}}{(b-a)t}$
$\sigma^2$	$\mu_r = 0, r$ odd; $\mu_r = \frac{r!}{(r/2)! 2^{r/2}}$ , $r$ even; $\kappa_r = 0, r > 2$	$\exp[\mu t + \frac{1}{2} \sigma^2 t^2]$
$\frac{1}{\lambda^2}$	$\mu_r' = \frac{\Gamma(r+1)}{\lambda^r}$	$\frac{\lambda}{\lambda-t}$ for $t < \lambda$
$\frac{r}{\lambda^2}$	$\mu_r' = \frac{\Gamma(r+1)}{\lambda^r \Gamma(r)}$	$\left(\frac{\lambda}{\lambda-t}\right)^r$ for $t < \lambda$
$\frac{ab}{(a+b+1)(a+b)^2}$	$\mu_r' = \frac{B(r+a, b)}{B(a, b)}$	not useful
Does not exist	Does not exist	Characteristic function is $e^{t\alpha - \beta t }$
$\exp[2\mu + 2\sigma^2] - \exp[2\mu + 2\sigma^2]$	$\mu_r' = \exp[r\mu + \frac{1}{2} r^2 \sigma^2]$	not useful
$2\beta^2$	$\mu_r = 0$ for $r$ odd; $\mu_r = r! \beta^r$ for $r$ even	$\frac{e^{at}}{1 - (\beta t)^2}$

Table 2 CONTINUOUS DISTRIBUTIONS (continued)

Name of parametric family of distributions	Cumulative distribution function $F(x)$ or probability density function $f(x)$	Parameter space	Mean $\mu = E\{X\}$
Weibull	$f(x) = abx^{b-1} \exp[-ax^b] I_{(0, \infty)}(x)$	$a > 0$ $b > 0$	$a^{-1/b} \Gamma(1 + b^{-1})$
Logistic	$F(x) = [1 + e^{-(x-\alpha)/\theta}]^{-1}$	$-\infty < \alpha < \infty$ $\theta > 0$	$\alpha$
Pareto	$f(x) = \frac{\theta x_0^\theta}{x^{\theta+1}} I_{(x_0, \infty)}(x)$	$x_0 > 0$ $\theta > 0$	$\frac{\theta x_0}{\theta - 1}$ for $\theta > 1$
Gumbel or extreme value	$F(x) = \exp(-e^{-(x-\alpha)/\theta})$	$-\infty < \alpha < \infty$ $\theta > 0$	$\alpha + \beta\gamma$ $\gamma \approx .577216$
$t$ distribution	$f(x) = \frac{\Gamma((k+1)/2)}{\Gamma(k/2)} \frac{1}{\sqrt{k\pi}} \frac{1}{(1+x^2/k)^{k+1/2}}$	$k > 0$	$\frac{\mu=0}{\text{for } k > 1}$
$F$ distribution	$f(x) = \frac{\Gamma((m+n)/2)}{\Gamma(m/2)\Gamma(n/2)} \left(\frac{m}{n}\right)^{m/2} \times \frac{x^{m-2}}{[1+(m/n)x]^{m+n/2}} I_{(0, \infty)}(x)$	$m, n = 1, 2, \dots$	$\frac{n}{n-2}$ for $n > 2$
Chi-square distribution	$f(x) = \frac{1}{\Gamma(k/2)} \left(\frac{1}{2}\right)^{k/2} x^{k/2-1} e^{-(1/2)x} I_{(0, \infty)}(x)$	$k = 1, 2, \dots$	$k$

  

Variance $\sigma^2 = E\{(X - \mu)^2\}$	Moments $\mu'_r = E\{X^r\}$ or $\mu'_r = E\{(X - \mu)^r\}$ and/or cumulants $k_r$	Moment generating function $e^{tX}$
$a^{-2/b} \frac{\Gamma(1 + 2b^{-1})}{\Gamma^2(1 + b^{-1})}$	$\mu'_2 = a^{-1/b} \Gamma\left(1 + \frac{2}{b}\right)$	$e^{tX} = a^{-1/b} \Gamma\left(1 + \frac{t}{b}\right)$
$\frac{\beta^2 \pi^2}{3}$		$e^{it\pi\beta t} \csc(\pi\beta t)$
$\frac{\theta x_0^2}{(\theta - 1)^2 (\theta - 2)}$ for $\theta > 2$	$\mu'_2 = \frac{\theta x_0^2}{\theta - 1}$ for $\theta > r$	does not exist
$\frac{\pi^2 \beta^2}{6}$	$k_2 = (-\beta\gamma\psi^{(r-1)}(1))$ for $r \geq 2$ , where $\psi^{(r)}(z)$ is digamma function	$e^{it\Gamma(1 - \beta t)}$ for $t < 1/\beta$
$\frac{k}{k-2}$ for $k > 2$	$\mu'_2 = 0$ for $k > r$ and $r$ odd $\mu'_2 = \frac{k^{r/2} B(r+1/2, (k-r)/2)}{B(k/2, k/2)}$ for $k > r$ and $r$ even	does not exist
$\frac{2n^2(m+n-2)}{m(n-2)^2(n-4)}$ for $n > 4$	$\mu'_2 = \left(\frac{n}{m}\right) \frac{\Gamma(m/2 + r) \Gamma(n/2 - r)}{\Gamma(m/2) \Gamma(n/2)}$ for $r < \frac{n}{2}$	does not exist
$2k$	$\mu'_2 = \frac{2\Gamma(k/2 + 1)}{\Gamma(k/2)}$	$\left(\frac{1}{1-2t}\right)^{k/2}$ for $t < 1/2$